

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
21 May 2004 (21.05.2004)

PCT

(10) International Publication Number
WO 2004/042350 A2


- (51) International Patent Classification⁷: G01N (74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).
- (21) International Application Number: PCT/US2003/033978 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 24 October 2003 (24.10.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/422,356 30 October 2002 (30.10.2002) US (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): GOKLEN, Kent, E. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). NTI-GYABAAH, Joseph [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). ANTIA, Firoz, D. [IN/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). DAHLGREN, Mary, Ellen [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: USING AMINES OR AMINO ACIDS AS MOBILE PHASE MODIFIERS IN CHROMATOGRAPHY



(57) Abstract: This invention relates to the use of amine, amino acid or amino acid ester mobile phase modifiers in normal phase chromatography to improve the resolution and/or productivity of peptide and lipopeptide purification. This chromatographic method can be used for either on analytical or preparative scale purification.

TITLE OF THE INVENTION

USING AMINES OR AMINO ACIDS AS MOBILE PHASE MODIFIERS IN
CHROMATOGRAPHY

5 CROSS REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application No. 60/422,356, filed October 30, 2002.

10 FIELD OF THE INVENTION

This invention relates to the use of amines, amino acids or amino acid esters as mobile phase modifiers in normal phase chromatography of peptide and lipopeptide compounds.

BACKGROUND OF THE INVENTION

15 In the past, the resolution of key impurities such as Pneumocandins B₅ and E₀ from Pneumocandin B₀ in the silica gel HPLC purification was poor. Some analogs were only partially resolved from the main product peak under preparative conditions. To achieve the desired product purity, the limited resolution required that the purification step be run at low column loading, which limited productivity.


Lipopeptides, such as Pneumocandin B₀, are often the product of a fermentation
20 process. During such a fermentation process, many very closely related analogues are produced along with the desired product. Normal phase chromatography systems are frequently used to purify the crude fermentation product. A normal phase chromatography system usually consists of a stationary phase and a mobile phase. For purification of a peptide or lipopeptide, the stationary phase can be silica gel or alumina, and the mobile phase can be a single solvent or a
25 mixture of solvents, which includes organic solvents and water.

Silica gel chromatography and other types of normal phase chromatography are useful for separating these analogues. However, in practice, the resolution of certain close analogues from the desired product is often poor and not satisfactory, because the resolution is not great and often there is overlap. To achieve the desired purity of the main product at a
30 reasonable yield requires restricting the amount of material (often referred to as feed or column load) loaded onto the column per run, which limits the productivity of the operation.

The purification of Pneumocandin B₀ falls into this category. The chromatography utilizes a mobile phase consisting of a mixture of solvents, specifically ethyl acetate (EtOAc), methanol (MeOH) and water, on a silica gel column. Pneumocandin B₀, with a
35 molecular weight of 1065 Daltons, is a natural product and serves as an intermediate in the

production of Caspofungin acetate (Cancidas®). Pneumocandin B₀ is produced as a secondary metabolite by fermentation of the fungus *Glarea lozoyensis*. See US Patent Nos. 5, 194,377 and 5,202,309. The structures of Pneumocandin B₀ and two of the key analog impurities, all comprised of a cyclic hexapeptide coupled with dimethylmyristate side chain, are shown in

5 Formula I and Table 1.



Formula I

Table 1. Pneumocandin B₀ and two of its analogs

Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶
Pneumocandin B ₀	OH	OH	Me	OH	H	OH
Pneumocandin B ₅	OH	H	Me	OH	H	OH
Pneumocandin E ₀	OH	OH	Me	OH	H	H

Silica gel chromatography exploits the subtle variations in binding affinity of the hydroxy-rich cyclic hexapeptide core of the desired product and the analog impurities, including Pneumocandins B₅ and E₀, to effect a separation. In the silica gel HPLC purification, Pneumocandins B₅ and E₀, two of the key analog impurities co-produced in the fermentation of Pneumocandin B₀, elute very close to Pneumocandin B₀. Therefore, to meet the target impurity levels in the purified material for these and similar analog compounds, the quantity of crude

20 Pneumocandin B₀ that can be loaded onto the column is limited. As a result, significant efforts have been made to improve the resolution of key impurities. For instance, the ternary ethyl

acetate-methanol-water mobile phase has been balanced to optimize resolution between Pneumocandin B₀ and key analog impurities. D.J. Roush, F.D. Antia, K.E. Göklen *J. Chromatography A*, 827 (1998) 373-389.

5 SUMMARY OF THE INVENTION

This invention discloses the use of mobile phase modifiers, including amino acids, amino acid esters or amines, during the HPLC purification of a lipopeptide or peptide. The mobile phase modifiers bind to the stationary phase and modify the binding characteristics of the stationary phase improving the resolution and or purification of desired lipopeptide or peptide
10 from its related impurities.

This invention is useful in analytical chromatography and even more valuable for preparative chromatography (i.e., using chromatography as a large-scale purification technique). Specifically, amino acids, amino acid esters or amines can be used as mobile phase modifiers in the chromatographic purification of peptides or lipopeptides, for example the purification of
15 Pneumocandin B₀, which is the natural product starting material (a fermentation product) used to prepare Caspofungin acetate (Cancidas®). The invention would also be useful in purifying the fermentation product precursor for other lipopeptides, such as Micafungin, Anidulafungin, Cilofungin and Daptomycin.

20 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1

Preparative HPLC chromatogram of silica gel chromatography of crude Pneumocandin B₀, where the column feed contains a small amount of proline. Figure 1A. First run on the column. Figure 1B. Eighteenth run on the column. Improved resolution of analogue impurities is seen in the
25 later run due to the adsorption of proline from the feed solution onto the column.

Figure 2

Figure 2A. Plot of retention time of Pneumocandin B₀ after injecting proline-containing solution onto the column, showing increased retention time, which returns to the original retention time, as the proline is slowly desorbed from the column. Figure 2B and 2C. Chromatogram of a
30 typical run just before (2B) and after (2C) injecting proline onto the column. (analytical scale 5μ YMC silica column using 87/9/7 v/v/v ethyl acetate/methanol/water).

Figure 3

Silica gel HPLC chromatograms for Pneumocandin B₀ crude eluted with mobile phase: (3A) without L-proline, (3B) with 0.26mM L-proline added, and (3C) with 0.65mM L-proline added. Demonstrates increased retention time of Pneumocandin B₀ and increased resolution from analog impurities with increasing levels of proline added. (Pneumocandin B₀ retention time is 22 minutes with 0.65mM L-proline).

Figure 4

Silica gel HPLC chromatogram for Pneumocandin B₀ crude eluted using 20% ethyl acetate and 80% 87/9/7 v/v/v ethyl acetate/ methanol/ water as the mobile phase. This mobile phase mixture results in increased retention time of Pneumocandin B₀, but without the improved resolution from its analogs observed when proline is added to the mobile phase (see Figure 3). (Pneumocandin B₀ retention time is 23 minutes).

Figures 5 through 11

Silica gel HPLC chromatograms for Pneumocandin B₀ crude eluted (A) without mobile phase modifier (control) and (B) with the addition of various amino acids to the mobile phase. Demonstrates that addition of all the amino acids have an affect on the Pneumocandin B₀ retention and/or its resolution from its analogs. (analytical scale 5 μ YMC silica column using 87/9/7 v/v/v ethyl acetate/methanol/water).

Figures 5

Figure 5A. Control prior to exposure of column to trans-4-hydroxyl-L-proline.

Figure 5B. Chromatogram after exposure of column to mobile phase modified with trans-4-hydroxyl-L-proline, showing similar retention time of Pneumocandin B₀ to the control, but different resolution from its analog impurities.

Figures 6

Figure 6A. Control prior to exposure of column to L-valine.

Figure 6B. Chromatogram after exposure of the column to mobile phase modified with L-valine, showing changes from the control in both the retention time of Pneumocandin B₀ and the resolution from its analog impurities.

Figures 7

Figure 7A. Control prior to exposure of column to L-lysine.

Figure 7B. Chromatogram after exposure of the column to mobile phase modified with L-lysine, showing changes from the control in both the retention time of Pneumocandin B₀ and the resolution from its analog impurities.

5 Figures 8

Figure 8A. Control prior to exposure of column to L-methionine.

Figure 8B. Chromatogram after exposure of the column to mobile phase modified with L-methionine, showing changes from the control in both the retention time of Pneumocandin B₀ and the resolution from its analog impurities.

10

Figures 9

Figure 9A. Control prior to exposure of column to D-proline.

Figure 9B. Chromatogram after exposure of the column to mobile phase modified with D-proline, showing changes from the control in both the retention time of Pneumocandin B₀ and the resolution from its analog impurities.

15

Figures 10

Figure 10A. Control prior to exposure of column to L-threonine.

Figure 10B. Chromatogram after exposure of the column to mobile phase modified with L-threonine, showing changes from the control in both the retention time of Pneumocandin B₀ and the resolution from its analog impurities.

20

Figures 11

Figure 11A. Control prior to exposure of column to glycine.

Figure 11B. Chromatogram after exposure of the column to mobile phase modified with glycine, showing changes from the control in both the retention time of Pneumocandin B₀ and the resolution from its analog impurities.

25

Figure 12

Silica gel HPLC chromatograms for Pneumocandin B₀ crude eluted (A) without (control) and (B) with the addition of diethylamine to the mobile phase. Demonstrates that diethylamine addition has a very strong affect on the Pneumocandin B₀ retention and its resolution from its analogs. (analytical scale 5 μ YMC silica column using 87/9/7 v/v/v ethyl acetate/methanol/water).

30

35 DETAILED DESCRIPTION OF THE INVENTION

A method for the purification of a peptide or a lipopeptide by using a mobile phase modifier in a normal phase chromatography system to improve the selectivity and/or productivity of the purification is disclosed. The method as recited above, wherein the mobile phase modifier is selected from a group consisting of an amine, an amino acid or an amino acid ester. The method as recited above, wherein the normal phase chromatography system includes a mobile phase and a stationary phase. The method as recited above, wherein the stationary phase is selected from silica gel and alumina. The method as recited above, wherein the mobile phase is a solvent system comprising one or more solvents. The method as recited above, wherein the mobile phase modifier is an amine. The method as recited above, wherein the amine is a primary, secondary, or tertiary amines, including lower alkyl (C₁-C₆ alkyl) amines, lower dialkyl (C₁-C₆ alkyl) amines and /or aromatic (C₆- C₁₀ aryl) amines. Examples of amines, which are useful in the method, as recited above are methylamine, ethylamine, diisopropylamine, diethylamine, dimethylamine, ethylmethylamine, triethylamine, propylamine, aniline and dimethylaniline. The method as recited above, wherein the mobile phase modifier is an amino acid or amino acid ester. The method as recited above, wherein the amino acid or amino acid ester mobile phase modifier is selected from the group consisting of: L-amino acids, D-amino acids, L-amino acid esters and D-amino acid esters. The method as recited above, wherein the amino acid or amino acid ester mobile phase modifier is selected from: L-proline, D-proline, *trans*-4-hydroxy-L-proline, *trans*-4-hydroxy-D-proline, glycine, L-threonine, D-threonine, L-lysine, D-lysine, L-methionine, D-methionine, D-valine, L-valine and esters of the aforementioned L-and D-amino acids. The method as recited above, wherein the amino acid is selected from: L-proline and D-proline.

An embodiment of the invention is the method of purifying a peptide or a lipopeptide by using an amine or amino acid mobile phase modifier in a normal phase chromatography system with a stationary phase of alumina or silica gel to improve the resolution and/or productivity of the purification, wherein the purification is of a peptide. Examples of peptides are linear and cyclic amino acid chains under 1800 molecular weight, formed by the combination of the amino group of one amino acid with the carboxyl group of another in an amide bond, and produced as secondary metabolites of microorganisms. Other examples of peptides are oxytocin and bradykinin.

A further embodiment of the invention is the method of purifying a peptide or a lipopeptide by using an amine or amino acid mobile phase modifier in a normal phase chromatography system with a stationary phase of alumina or silica gel to improve the resolution and/or productivity of the purification, wherein the purification is of a lipopeptide. The method

as recited above, wherein the lipopeptide is selected from a fermentation product precursor of Caspofungin, Micafungin, Anidulafungin, Cilofungin, and Daptomycin. The method as recited above, wherein the lipopeptide is pneumocandin B₀. The method as recited above, wherein the mobile phase modifier is an amine. The method as recited above, wherein the amine mobile phase modifier is selected from the group consisting of: methylamine, ethylamine, diisopropylamine, diethylamine, dimethylamine, ethylmethylamine, triethylamine, propylamine, aniline and dimethylaniline. The method as recited above, wherein the mobile phase modifier is an amino acid or amino acid ester. The method as recited above, wherein the amino acid or amino acid ester mobile phase modifier is selected from the group consisting of: L-amino acids, D-amino acids, L-amino acid esters and D-amino acid esters. The method as recited above, wherein the stationary phase is silica gel. The method as recited above, wherein the amino acid or amino acid ester mobile phase modifier is selected from: L-proline, D-proline, *trans*-4-hydroxy-L-proline, *trans*-4-hydroxy-D-proline, glycine, L-threonine, D-threonine, L-lysine, D-lysine, L-methionine, D-methionine, D-valine, L-valine and esters of the aforementioned L- and D-amino acids. The method as recited, wherein the mobile phase is a solvent system comprising water, methanol, and ethyl acetate. The method as recited above, wherein the amino acid mobile phase modifier is selected from: L-proline and D-proline.

Yet another embodiment of the invention is the method of purifying a peptide or a lipopeptide by using an amine or amino acid mobile phase modifier in a normal phase chromatography system with a stationary phase of alumina or silica gel to improve the resolution and/or productivity of the purification, wherein the purification is of a peptide. The method as recited above, wherein the mobile phase modifier is an amine. The method as recited above, wherein the mobile phase modifier is an amino acid or amino acid ester. The method as above, wherein the stationary phase is silica gel.

Examples of lipopeptides, for which this purification process is useful, are echinocandin derivatives, such as Pneumocandin B₀, Caspofungin, Cilofungin and Micafungin as well as Anidulafungin and Daptomycin, and particularly the natural product precursors of Caspofungin, Micafungin, Cilofungin, Anidulafungin and Daptomycin. The natural product/fermentation product precursor for Caspofungin is Pneumocandin B₀. Caspofungin acetate (CANCIDAS) is a semisynthetic lipopeptide echinocandin B derivative currently being sold in the US as an antifungal agent for intravenous administration. Anidulafungin is a semisynthetic lipopeptide echinocandin B derivative under development by Eli Lilly/Versicor as an antifungal agent for intravenous administration. Anidulafungin is disclosed in US Patent Nos. 5,965,525 and 6,384,013, hereby incorporated by reference. Cilofungin is an echinocandin lipopeptide disclosed by Eli Lilly in US Patent No. 4,293,489 for use as an antifungal agent, hereby

incorporated by reference. Micafungin (FUNGARD) is an echinocandin-like lipopeptide under development by Fujisawa, as an antifungal agent for intravenous administration. Micafungin is disclosed in US Patent No. 6,107,458 hereby incorporated by reference. Daptomycin (CIDEVIN) is a semisynthetic lipopeptide derivative under development by Cubist as an antibacterial agent. Daptomycin is disclosed by Eli Lilly in US Patent No. 4,537,717 hereby incorporated by reference.

A normal phase chromatography system employs a mobile phase and a stationary phase. The mobile phase is a solvent system comprising one or more solvents, the composition of which is either constant through out the purification process, or a gradient, where the solvent composition is changed over time during the purification process. The mobile phase solvents include, but are not limited to, water, methanol, ethanol, isopropanol, hexane, ethyl acetate, acetonitrile, and methylene chloride. The stationary phase is selected from silica gel and alumina. The instant invention provides a chromatographic purification method for a peptide or lipopeptide, which employs the addition of a mobile phase modifier to the mobile phase. A column volume is defined, as the volume of solvent needed to traverse the column. Column load (also referred to as column feed or feed load) refers to the amount of material (crude lipopeptide or peptide) that can be purified at one time. The mobile phase modifier is defined as an amino acid, an amino acid ester, or amine. The instant invention contemplates the addition of the mobile phase modifier either to the eluant (mobile phase solvent system) or as a supplement to the column load. Examples of the amino acids contemplated by the invention are any natural or unnatural amino acids, including both L- and D-configurations. Embodiments of the amino acid mobile phase modifiers useful in this process are L- and D-proline, *trans*-4-hydroxy-L-proline, glycine, L-threonine, L-lysine, L-methionine and L-valine. Examples of amine mobile phase modifiers contemplated by the invention are primary, secondary, or tertiary lower alkyl (C₁-C₆ alkyl) and /or aromatic (C₆- C₁₀ aryl) amines. Examples of such amines are methylamine, ethylamine, diisopropylamine, diethylamine, dimethylamine, ethylmethylamine, triethylamine, propylamine, aniline and dimethylaniline.

This invention shows that adding low concentrations of certain amines, amino acids, or amino acid esters to the mobile phase can shift the resolution and selectivity of the system, providing significant enhancement. In particular, the addition of proline to the EtOAc-MeOH-water mixture used in the silica gel chromatography of Pneumocandin B₀ significantly increases the resolution of the compound from closely related impurities, including Pneumocandins B₅ and E₀. This increased resolution allows the loading of the product onto the column to be significantly increased, by a factor of 2 to 3-fold, while maintaining the purity of the product rich cut and the yield. Larger amounts of product can be purified to the same purity

at the same yield on the same column of silica using less solvent and in less time. It has also been demonstrated that the presence of the mobile phase modifier in the column load will have a similar effect.

5 Solvents purchased from Fisher Scientific (Pittsburgh, PA, USA) were used. Analytical scale HPLC columns (250 x 4.6 mm id) packed with regular bare silica, 5 μ m particle size and 120 Å pore diameter, were obtained from YMC (Wilmington, NC, USA). The semi-preparative scale experiment employed irregular silica from Amicon (Beverly, MA, USA) designated Grade 631, Si-60 of nominal particle size 20 μ m. For analytical scale experiments, all
10 solvents in the ternary mobile phase and the feed diluent (ethyl acetate, methanol and water) were HPLC grade. Mobile phase employed for the semi-preparative experiment was prepared from methanol (99.9% w/w purity) from Enron (Houston, TX, USA) and urethane grade ethyl acetate from Eastman (Perth Amboy, NJ, USA) and low ion water produced in the laboratory using a purification unit from Osmonics (Minnetonka, MN, USA). For both the analytical and semi-prep
15 scale experiments, the mobile phase composition was 87/9/7 (v/v/v) ethyl acetate/methanol /water. Volumetric compositions of solutions do not account for non-ideal mixing effects. For the sake of convenience, the mobile phase will be referred to as WEAM (water/ethyl acetate/methanol). The feed used for the experiments contained Pneumocandin B₀ dissolved in a solvent blend of 79.3/13.3/7.4, v/v/v ethyl acetate/methanol/water. L-proline was obtained from
20 Kyowa Hakko Kogyo Ltd. (Japan). Glycine, L-threonine, D-proline, L-methionine, L-lysine, L-valine, trans-4-hydroxy-L-proline, diethyl amine and proline methyl ester were obtained from Sigma-Aldrich Cheme GmbH (Steinheim, Germany).

Analytical scale experiments were performed on a Hewlett Packard (Waldbronn, Germany) HP1100 HPLC system with column thermostat and variable wavelength detector
25 (VWD) with detection monitored at 278 nm. Semi-preparative experiments employed a Dorr Oliver Model C (1 LPM pumping skid, Biotage Division of Dyax) and a 50 cm x 6 cm id axial compression column from Prochrom (Indianapolis, IN).

The modifier effect was first observed during the purification of a particular batch of crude Pneumocandin B₀ product containing L-proline, which had been added into the
30 fermentation mixture to facilitate the fermentation process.

When the crude fermentation product of Pneumocandin B₀ was first loaded to the HPLC, the resolution of Pneumocandin B₀ was limited. Pneumocandin E₀ was buried under the peak for Pneumocandin B₀ and the separation of Pneumocandin B₅ from Pneumocandin B₀ was not prominent. See Figure 1A, which represents the UV (278 nm) traces for the 1st run of the
35 silica gel HPLC.

However, after a few sequential HPLC purification runs of this Pneumocandin B₀ fermentation product, an increase in the retention time of Pneumocandin B₀ by as much as 10 minutes was observed. In addition, Pneumocandin E₀, the Pneumocandin B₀ that co-eluted in earlier runs was now eluting ahead of the main peak. Furthermore, the resolution of
5 Pneumocandin B₅, the close eluter of Pneumocandin B₀, improved dramatically. Figure 1B presents the 18th in this series of HPLC runs and clearly shows this increase in retention time and enhanced resolution.

The enhanced separation was reversed by a methanol wash of the column, which was performed every twenty runs for the purpose of removing tightly bound impurities. The
10 improved purification results returned again after a few additional runs. Further investigation determined that the effect was due to the presence of proline in the column feed (the crude Pneumocandin B₀), which adsorbed to the column with each injection. This phenomenon was generalized by addition of proline and related additives to the mobile phase so that the level of the effect could be controlled, providing improvements in resolution and productivity, as
15 described in the examples.

The examples provided herein are intended to assist in a further understanding of the invention. Particular materials, employed species and conditions are intended to be further illustrative of the invention and not limitative of the reasonable scope thereof.

20 Example 1

On an analytical scale YMC 5 μ silica column using 87/9/7 v/v/v ethyl acetate/methanol/water as the mobile phase, a series of 8 injections of Pneumocandin B₀ product containing no L-proline were carried out to establish an unchanging control for comparison. Following this, 10 short injections of an L-proline solution (1.3×10^{-2} M L-proline in
25 79.3/13.3/7.4 v/v/v ethyl acetate/methanol/water, 60 μ L injection, about 5 min run time at 1.2 mL/min) were made, and nine additional injections of the Pneumocandin B₀ product containing no L-proline were carried out. The procedure was repeated if necessary. Figure 2A shows the retention time for the Pneumocandin B₀ peak in the system for injections made immediately after the series of proline injections, relative to the retention time prior to proline injection. It can be
30 seen that there was a marked increase in retention immediately after L-proline treatment. This was accompanied by an improvement in the selectivity between Pneumocandin B₀ and its immediate early eluting peak. However, the retention time returned to normal (RRT~1) after about 5 runs and the improved selectivity was lost, presumably due to the desorption of L-proline with continued elution from the column with the proline-free mobile phase.

Figures 2B and 2C are typical chromatograms obtained from the injections just before and just after the L-proline treatment. The change in retention is apparent, as is a distinct change in resolution between Pneumocandin B₀ and Pneumocandins B₅ and E₀.

5

Example 2

Experiments were carried out on a 5 Tm YMC silica-packed analytical column with L-proline added directly to the mobile phase. Figures 3A, B and C shows chromatograms without L-proline and with 0.26 mM (~ 30 mg/L) and 0.65 mM (~ 75 mg/L) L-proline in the mobile phase, respectively. The column was equilibrated for 2 hours at 1.2 ml/min with the proline containing mobile phases before the injections were carried out. In Figure 3C, the peak just ahead of Pneumocandin B₀ was determined to be Pneumocandin E₀ by the injection of standards. Pneumocandin E₀ was not separable when the chromatography was carried out without L-proline.

15

Example 3

The selectivity enhancement was studied by increasing the retention time of Pneumocandin B₀ to around 23 minutes, which is comparable to the retention time of around 22 minutes seen when 0.65 mM L-proline was used in Example 2, by eluting with a mobile phase of 20% ethyl acetate and 80% of the 87/9/7 v/v/v ethyl acetate/methanol/water. Although the retention time of Pneumocandin B₀ was increased, Pneumocandin E₀ was still buried in the peak of Pneumocandin B₀. No enhanced resolution was achieved. See Figure 4.

20

25

Example 4

On a semi-preparative scale, a 6 cm ID Prochrom column was packed with 390 g of Amicon Grade 631, 18-20 µ, 60 Å silica (slurried in a 87/9/7 v/v/v water /ethyl acetate / methanol (WEAM) mobile phase; packing pressure 40 bar; column length 25 cm) was used.

Benchmark Run:

A feed solution was prepared by dissolving 90g of crude Pneumocandin B₀ without L-proline in 1.5 L feed solvent (79.3/13.3/7.4 v/v/v ethyl acetate/methanol/water) giving a 34.9 assay g/L Pneumocandin B₀ feed solution. In order to establish a control for the experiment, 80 mL of the feed (2.8 g Pneumocandin B₀) were injected and eluted with the WEAM mobile phase at a flow rate of 90 mL/min. Fractions were collected from this run and analytical results of a pooled representative rich cut indicated that the impurity profile was

30

35

consistent with previous batches (by both normal phase (NP) and reversed phase (RP) HPLC assays).

Runs with L-proline in the mobile phase(similar load to benchmark):

- 5 The WEAM mobile phase was modified with L-proline at a concentration of 75 mg/L (0.650 mM). The column was conditioned with 5 column volumes (CV) of this solution and a run was carried out with the modified mobile phase under conditions otherwise similar to the benchmark runs. The shift in retention time observed in this system was not as dramatic as in the analytical system. Table 2 summarizes the impurity profiles of the rich cuts from the two runs: levels of Pneumocandin B₅ and Pneumocandin E₀ are dramatically lower in the proline modified run, showing that higher throughputs could be achieved with this system.
- 10

Table 2: Summary of rich cut impurity profiles from laboratory runs, standard loading, without and with 0.15 g/L (1.3 M) of L-proline in the mobile phase

Run	Load as grams Pneumocandin B ₀ per kg silica	Pn B ₅ % Area	Pn E ₀ % Area	Yield %	Mass Balance %
Benchmark run	7.2	0.44	0.42	82	99
L-proline system	7.2	0.16	0.05	84	98

15 *Doubling column throughput with the L-proline modified mobile phase:*

- A run was carried out using the proline-modified system described above with double the benchmark load: 5.6 g Pneumocandin B₀ injected (14.4 g/kg silica) by doubling the volume of the feed solution injected, all other conditions being the same. The impurity profile of the rich cut, as well as the yield and mass balance, is summarized in Table 3. The results indicate the potential use of this method to double the productivity of the chromatography step, since in this example the yield has been maintained while increasing purity at double the benchmark loading.
- 20

Table 3: Summary of rich cut impurity profiles from laboratory runs, 2X loading, without and with 0.075 g/L (0.65M) and 0.15 g/L (1.3 M) L-proline in the mobile phase

Run	Load: g Pneumocandin B ₀ per kg silica	Pn B ₅ % Area	Pn E ₀ % Area	Yield (%)	Mass Balance (%)
Benchmark	7.2	0.44	0.42	82	99
L-proline system	7.2	0.16	0.05	84	98
L-proline system	14.4	0.27	0.08	86	99

Example 6

Other amino acids as mobile phase modifiers

In order to obtain further insight regarding the phenomenon, a series of analytical experiments were carried out using a 25 x 0.46 cm YMC 5 μ silica column with a wide variety of amino acid modifiers, including *trans*-4-hydroxy-L-proline, L-valine, L-lysine, L-methionine, D-proline, L-threonine and glycine in the mobile phase at a concentration of 0.65 mM. See Figures 5-11.

Table 4 sorts the amino acids in order of decreasing retention. Figures 5 through 10 provide exemplary chromatograms obtained from these runs that illustrate their various effects on retention time and selectivity. With every amino acid except 4-hydroxy-L-proline, retention increased compared to the amino acid-free case, and the selectivity was modified uniquely in each instance. For example, although *trans*-4-hydroxy-L-proline did not cause the retention of Pneumocandin B₀ to shift, it did affect the resolution of Pneumocandin B₀ from the early eluting peak, as seen in Figure 5.

Glycine has no side chain, but it also increases the retention time. This strongly suggests that the retention time shift and the improved resolution is not the unique effect of the amino acid side chain.

L-proline methyl ester was employed, in place of an amino acid and had an increased retention time (retention time of Pneumocandin B₀ relative to the control was 1.65) and a change in selectivity.

After each set of 3 or 4 runs with a particular amino acid, the column was washed with 36 ml (~8 column volumes, hereinafter referred to as CV) of methanol and then re-equilibrated with 10 CV of neat WEAM mobile phase, after which three (~30 min) injections

were carried out to confirm that the amino acid had been removed (i.e., that baseline performance had been recovered). Following this, the column was re-conditioned with 72 mL (~17 CV) of the mobile phase containing a new amino acid, and new injections were then made. The retention times reported are for the 3rd run in each case. Since it is unclear what the saturation capacity of the silica is for the different amino acids, this protocol ensured that in each case the column had been exposed to the same number of moles of the amino acid. The column could not be adequately regenerated after exposure to L-lysine (which was purposely tested as the last amino acid), presumably due to the two primary amines.

Table 4. Retention times of Pneumocandin B₀ on a YMC 5 μ , 120 Å, silica column
25 cm x 0.46 cm ID at a flow rate of 1.2 mL/min.

Amino Acid (65 mM) in WEAM	Retention of Pneumocandin B ₀ relative to the control
L-proline methyl ester	1.65
L-proline	1.44
Glycine	1.38
L-threonine	1.35
D-proline	1.25
L-methionine	1.16
L-lysine	1.13
L-valine	1.10
<i>trans</i> -4-hydroxy-L-proline	1.00
None (control)	1.00

Example 7

Chromatography with Diethylamine as a mobile phase modifier

Diethylamine was added to the mobile phase at about 0.1 g/L, and the experimental conditions were identical to that employed previously in Example 6. As shown in Figure 12, the retention time increased even more with diethylamine than with proline. Therefore, the influence on retention and selectivity appears to be mediated by the amine functionality, such that either an amine or an amino acid/ amino acid ester could be used as a mobile phase modifier in this system.

WHAT IS CLAIMED IS:

1. A method of purifying a peptide or a lipopeptide by using a mobile phase modifier in a normal phase chromatography system to improve the selectivity and/or productivity of the purification.
2. The method as recited in claim 1, wherein the mobile phase modifier is selected from a group consisting of an amine, an amino acid or an amino acid ester.
3. The method as recited in claim 2, wherein the normal phase chromatography system includes a mobile phase and a stationary phase.
4. The method as recited in claim 3, wherein the stationary phase is selected from silica gel and alumina.
5. The method as recited in claim 4, wherein the mobile phase is a solvent system comprising one or more solvents.
6. The method as recited in claim 5, wherein the mobile phase modifier is an amine.
7. The method as recited in Claim 6, wherein the amine is selected from the group consisting of: methylamine, ethylamine, diisopropylamine, diethylamine, dimethylamine, ethylmethylamine, triethylamine, propylamine, aniline and dimethylaniline.
8. The method as recited in claim 5, wherein the mobile phase modifier is an amino acid or amino acid ester.
9. The method as recited in Claim 8, wherein the amino acid or amino acid ester mobile phase modifier is selected from the group consisting of: L-amino acids, D-amino acids, L-amino acid esters and D-amino acid esters.
10. The method as recited in Claim 9, wherein the amino acid or amino acid ester mobile phase modifier is selected from: L-proline, D-proline, *trans*-4-hydroxy-L-proline,

trans-4-hydroxy-D-proline, glycine, L-threonine, D-threonine, L-lysine, D-lysine, L-methionine, D-methionine, D-valine, L-valine and esters of the aforementioned L- and D-amino acids.

5 11. The method as recited in claim 10, wherein the amino acid is selected from: L-proline and D-proline.

12. The method as recited in claim 5, wherein the normal phase chromatography system is for the purification of a peptide.

10 13. The method as recited in claim 5, wherein the normal phase chromatography system is for the purification of a lipopeptide.

14. The method as recited in claim 13, wherein the lipopeptide is a fermentation product precursor of caspofungin, micafungin, cilofungin, andulifungin and
15 daptomycin.

15. The method as recited in claim 14, wherein the lipopeptide is a pneumocandin B₀.

20 16. The method as recited in claim 15, wherein the mobile phase modifier is an amine.

17. The method as recited in Claim 16, wherein the amine mobile phase modifier is selected from the group consisting of: methylamine, ethylamine, diisopropylamine, diethylamine, dimethylamine, ethylmethylamine, triethylamine, propylamine, aniline and
25 dimethylaniline.

18. The method as recited in claim 15, wherein the mobile phase modifier is an amino acid or amino acid ester.

30 19. The method as recited in Claim 18, wherein the amino acid or amino acid ester mobile phase modifier is selected from the group consisting of: L-amino acids, D-amino acids, L-amino acid esters and D-amino acid esters.

35 20. The method as in claim 19, wherein the stationary phase is silica gel.

21. The method as recited in Claim 20, wherein the amino acid or amino acid ester mobile phase modifier is selected from: L-proline, D-proline, *trans*-4-hydroxy-L-proline, *trans*-4-hydroxy-D-proline, glycine, L-threonine, D-threonine, L-lysine, D-lysine, L-methionine, D-methionine, D-valine, L-valine and esters of the aforementioned L- and D-amino acids.

22. The method as recited in claim 21, wherein the mobile phase is a solvent system comprising water, methanol, and ethyl acetate.

23. The method as recited in claim 22, wherein the amino acid mobile phase modifier is selected from: L-proline and D-proline.


24. The method as recited in claim 12, wherein the peptide is oxytocin or bradykinin.

25. The method as recited in claim 24, wherein the mobile phase modifier is an amine.


26. The method as recited in claim 24, wherein the mobile phase modifier is an amino acid or amino acid ester.

27. The method as in claim 26, wherein the stationary phase is silica gel.


28. A method of purifying a peptide or a lipopeptide by using a mobile phase modifier in a normal phase chromatography system to improve the selectivity and/or productivity of the purification, except that when the lipopeptide is Pneumocandin B₀, then the mobile phase modifier is not L-proline.




20959PV, Figure 1A




20959PV, Figure 1B




20959PV, Figure 2A




20959PV, Figure 2B




20959PV, Figure 2C



20959PV, Figure 3A




20959PV, Figure 3B




20959PV, Figure 3C


4/12




20959PV, Figure 4




20959PV, Figure 5A




20959PV, Figure 5B




20959PV, Figure 6A




20959PV, Figure 6B




20959PV, Figure 7A



20959PV, Figure 7B




20959PV, Figure 8A




20959PV, Figure 8B


9/12




20959PV, Figure 9A




20959PV, Figure 9B




20959PV, Figure 10A




20959PV, Figure 10B




20959PV, Figure 11A



20959PV, Figure 11B



20959PV, Figure 12A



20959PV, Figure 12B

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
21 May 2004 (21.05.2004)

PCT

(10) International Publication Number
WO 2004/042350 A3

(51) International Patent Classification⁷: **B01D 15/00**,
15/08, C07K 14/00

(21) International Application Number:
PCT/US2003/033978

(22) International Filing Date: 24 October 2003 (24.10.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/422,356 30 October 2002 (30.10.2002) US

(71) Applicant (*for all designated States except US*): **MERCK & CO., INC.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **GOKLEN, Kent, E.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **NTI-GYABAAH, Joseph** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **ANTIA, Firoz, D.** [IN/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **DAHLGREN, Mary, Ellen** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(74) Common Representative: **MERCK & CO., INC.**; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

(88) Date of publication of the international search report:
15 July 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: USING AMINES OR AMINO ACIDS AS MOBILE PHASE MODIFIERS IN CHROMATOGRAPHY

(57) Abstract: This invention relates to the use of amine, amino acid or amino acid ester mobile phase modifiers in normal phase chromatography to improve the resolution and/or productivity of peptide and lipopeptide purification. This chromatographic method can be used for either on analytical or preparative scale purification.



WO 2004/042350 A3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/33978

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : B01D 15/00, 15/08; C07K 14/00

US CL : 210/656, 660; 530/300, 350, 359, 412, 415, 417

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 210/656, 660; 530/300, 350, 359, 412, 415, 417

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4,784,988 (BRUZZESE et al) 15 November 1988 (15.11.1988), column 12, lines 25-68.	1-6 and 12
X	US 4,812,442 (BOGER et al) 14 March 1989 (15.11.1988), column 38, lines 34-50.	1-6 and 12
X	JP 11023558A (TOSOH CORP.) 29 January 1999 (29.01.1999) Abstract.	1-3
Y	US 6,153,590 A (ANDERSEN et al) 28 November 2000 (28.11.2000), column 24, lines 7-18.	1, 2, 4 and 5

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

13 March 2004 (13.03.2004)

Date of mailing of the international search report

19 MAY 2004

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

Facsimile No. (703)305-3230

Authorized officer

Valerie Bell-Harris
Chin-Min Kam

Telephone No. 571/272-1600

INTERNATIONAL SEARCH REPORT

PCT/US03/33578

Continuation of B. FIELDS SEARCHED Item 3:

EAST Search on USPAT, DERWENT, EPO, JPO, USPGPUB; STN search on MEDLINE, CAPLUS, BIOSIS, EMBASE, SCISEARCH, AGRICOLA. Search term used: chromatograph?, silica gel, alumina, normal phase, modifier, purif?, isolat?, amine, amino acid, aniline, peptide, lipopeptide, oxytocin, bradykinin, caspofungin, micafungin, cilogungin, andulifungin, daptomycin, pneumocandin, mobile phase, elut?.